

## **MAS References:**

Akman MN, Bengi R, Karatas M, Kiliç S, Sözü S, Ozker R. Assessment of spasticity using isokinetic dynamometry in patients with spinal cord injury. *Spinal Cord* 1999;Sep; 37(9):638-43.

Akpinar P, Atici A, Ozkan FU, Aktas I, Kulcu DG, Sari A, Durmus B. Reliability of the Modified Ashworth Scale and Modified Tardieu Scale in patients with spinal cord injuries. *Spinal Cord* 2017;55(10):944-949.

Alhusaini AA, Dean CM, Crosbie J, Shepherd RB, Lewis J. Evaluation of spasticity in children with cerebral palsy using Ashworth and Tardieu Scales compared with laboratory measures. *Journal of Child Neurology* 2010;25(10):1242-7.

Alibiglou L, Rymer WZ, Harvey RL, Mirbagheri MM. The relation between Ashworth scores and neuromechanical measurements of spasticity following stroke. *Journal of NeuroEngineering and Rehabilitation* 2008;5:18.

Allison SC, Abraham LD. Correlation of quantitative measures with the modified Ashworth scale in the assessment of plantar flexor spasticity in patients with traumatic brain injury. *Journal of Neurology* 1995;242(10):699-706.

Allison SC, Abraham LD. Sensitivity of qualitative and quantitative spasticity measures to clinical treatment with cryotherapy. *International Journal of Rehabilitation Research* 2001;24(1):15-24.

Allison SC, Abraham LD, Peterson CL. Reliability of the Modified Ashworth Scale in the assessment of plantarflexor muscle spasticity in patients with traumatic brain injury. *International Journal of Rehabilitation Research* 1996;19(1):67-78.

Annaswamy T, Mallempati S, Allison SC, Abraham LD. Measurement of plantarflexor spasticity in traumatic brain injury: correlational study of resistance torque compared with the modified Ashworth scale. *American Journal of Physical Medicine and Rehabilitation* 2007;86(5):404-11.

Ansari NN, Naghdi S, Arab TK, Jalaie S. The interrater and intrarater reliability of the Modified Ashworth Scale in the assessment of muscle spasticity: limb and muscle group effect. *NeuroRehabilitation* 2008;23(3):231-7.

Ansari NN, Naghdi S, Moammeri H, Jalaie SH. Ashworth Scales are unreliable for the assessment of muscle spasticity. *Physiotherapy Theory and Practice* 2006;22(3): 119-25.

Ashworth B. Preliminary trial of carisoprodol in multiple sclerosis. *Practitioner* 1964;192:540-2.

Bakheit AMO, Maynard VA, Curnow J, Hudson N, Kodapala S. The relation between Ashworth Scale scores and the excitability of the alpha motor neurons in patients with post-stroke muscle spasticity. *Journal of Neurology, Neurosurgery, and Psychiatry* 2003;74:646-648.

Bar-On L, Van Campenhout A, Desloovere K, Aertbelien E, Huenaeerts C, Vandendoorent B, Nieuwenhuys A, Molenaers G. Is an instrumented spasticity assessment an improvement over clinical spasticity scales in assessing and predicting the response to integrated botulinum toxin type a treatment in children with cerebral palsy? *Archives of Physical Medicine and Rehabilitation* 2014;95(3);515-23.

Biering-Sorensen F, Nielsen JB, Klinge K. Spasticity-assessment: a review. *Spinal Cord* 2006;44(12):708-722.

Boyd RN, Barwood SA, Baillieu Ce, Graham HK. Validity of a clinical measure of spasticity in children with cerebral palsy in a randomized clinical trial. *Developmental Medicine and Child Neurology* 1998;40(S78):7.

Boyd RN, Graham HK. Objective measurement of clinical findings in the use of botulinum toxin type A for the management of children with cerebral palsy. *European Journal of Neurology* 1999;6(S4):s23-35.

Blackburn M, van Vliet P, Mockett SP. Reliability of measurements obtained with the Modified Ashworth Scale in the lower extremities of people with stroke. *Physical Therapy* 2002;82(1):25-34.

Bodin PG, Morris ME. Inter-rater reliability of the Modified Ashworth Scale for wrist flexor spasticity following stroke. *Proceedings of the 11<sup>th</sup> Congress of the World Confederation of Physical Therapy* 1991:505-7.

Bohannon RW, Smith MB. Interrater reliability of a Modified Ashworth Scale of muscle spasticity. *Physical Therapy* 1987;67:206-7.

Clopton N, Dutton J, Featherston T, Grigsby A, Mobley J, Melvin J. Interrater and intrarater reliability of the Modified Ashworth Scale in children with hypertonia. *Pediatric Physical Therapy* 2005;17(4);268-74.

Cooper A, Musa IM, van Deursen R, Wiles CM. Electromyography characterization of stretch responses in hemiparetic stroke patients and their relationship with the Modified Ashworth scale. *Clinical Rehabilitation* 2005;19(7):760-6.

Craven BC, Morris AR. Modified Ashworth Scale reliability for measurement of lower extremity spasticity among patients with SCI. *Spinal Cord* 2010;48(3):207-13.

Damiano DL, Quinlivan JM, Owen BF, Payne P, Nelson KC, Abel MF. What does the Ashworth scale really measure and are instrumented measures more valid and precise? *Developmental Medicine and Child Neurology* 2002;44(2):112-8.

Fosang AL, Galea MP, McCoy AT, Reddihough DS. Measures of muscle and joint performance in the lower limb of children with cerebral palsy. *Developmental Medicine & Child Neurology* 2003;45(10):664-670.

Franzoi AC, Castro C, Cardone C. Isokinetic assessment of spasticity in subjects with traumatic spinal cord injury (ASIA A). *Spinal Cord* 1999Jun;37(6):416-20.

Gregson JM, Leathley M, Moore AP, Sharma AK, Smith TL, Watkins CL. Reliability of the Tone Assessment Scale and the Modified Ashworth Scale as clinical tools for assessing poststroke spasticity. *Archives of Physical Medicine and Rehabilitation* 1999;80(9):1013-1016.

Gregson JM, Leathley MJ, Moore AP, Smith TL, Sharma AK, Watkins CL. Reliability of measurements of muscle tone and muscle power in stroke patients. *Age and Ageing* 2000;29(3):223-8.

Haas BM, Bergström, Jamous A, Bennie A. The inter rater reliability of the original and of the Modified Ashworth Scale for the assessment of spasticity in patients with spinal cord injury. *Spinal Cord* 1996;34(9):560-4.

Heidari M, Abootalebi S, Hosseini SA. Validity of Modified Ashworth Scale as a measure of wrist spasticity in stroke patients. *Iranian Rehabilitation Journal* 2011;9(13):26-30.

Kaya T, Karatepe AG, Gunaydin R, Koc A, Altundal Ercan U. Inter-rater reliability of the Modified Ashworth Scale and Modified Modified Ashworth Scale in assessing poststroke elbow flexor spasticity. *International Journal of Rehabilitation Research* 2011;34(1):59-64.

Katz RT, Rovai GP, Brait C, Rymer WZ. Objective quantification of spastic hypertonia: correlation with clinical findings. *Archives of Physical Medicine and Rehabilitation* 1992;73(4):339-47.

Klingels K, De Cock P, Molenaers G, Desloovere K, Huenaerts C, Jaspers E, Feys H. Upper limb motor and sensory impairments in children with hemiplegic cerebral palsy. Can they be measured reliably? *Disability and Rehabilitation* 2010;32(5):409-16.

Kumar RTS, Pandyan AD, Sharma AK. Biomechanical measurement of post-stroke spasticity. *Age and Ageing* 2006;35:371-375.

Li F, Wu Y, Li X. Test-retest reliability and inter-rater reliability of the Modified Tardieu Scale and the Modified Ashworth Scale in hemiplegic patients with stroke. *European Journal of Physical and Rehabilitation Medicine* 2014;50(1):9-15.

Lin FM, Sabbahi M. Correlation of spasticity with hyperactive stretch reflexes and motor dysfunction in hemiplegia. *Archives of Physical Medicine and Rehabilitation* 1999;80(5): 526-530.

Mackey AH, Walt SE, Lobb G, Stott NS. Intraobserver reliability of the modified Tardieu scale in the upper limb of children with hemiplegia. *Developmental Medicine and Child Neurology* 2004;46(4):267-72.

Malhotra S, Cousins E, Ward A, Day C, Jones P, Roffe C, Pandyan A. An investigation into the agreement between clinical, biomechanical and neurophysiological measures of spasticity. *Clinical Rehabilitation* 2008;22(12):1105-1115.

Mehrholz J, Wagner K, Meißner D, Grundmann K, Zange C, Koch R, Pohl M. Reliability of the Modified Tardieu Scale and the Modified Ashworth Scale in adult patients with severe brain injury: a comparison study. *Clinical Rehabilitation* 2005;19(7):751-9.

Meseguer-Henarejos A-B, Sanchez-Meca J, Lopez-Pina J-A, Carles-Hernandez R. Inter- and intra-rater reliability of the Modified Ashworth Scale: a systemic review and meta-analysis. *European Journal of Physical Rehabilitation Medicine* 2017;Sep 13;(). doi: [10.23736/S1973-9087.17.04796-7](https://doi.org/10.23736/S1973-9087.17.04796-7) [Epub ahead of print] PMID: 28901119

Min JH, Shin YI, Joa KL, Ko SH, Shin MJ, Chang JH, Ko HY. The correlation between Modified Ashworth Scale and biceps T-reflex and inter-rater and intra-rater reliability of biceps T-reflex. *Annals of Rehabilitation Medicine* 2012;36(4): 538-543.

Mutlu A, Livanelioglu A, Gunel MK. Reliability of Ashworth and Modified Ashworth Scales in children with spastic cerebral palsy. *BMC Musculoskeletal Disorders* 2008;9:44.

Numanoğlu A, Günel MK. Intraobserver reliability of Modified Ashworth Scale and Modified Tardieu Scale in the assessment of spasticity in children with cerebral palsy. *Acta Orthopaedica et Traumatologica Turc* 2012;46(3):196-200.

Pandyan AD, Johnson GR, Price CI, Curless RH, Barnes MP, Rodgers H. A review of the properties and limitations of the Ashworth and Modified Ashworth Scales as measures of spasticity. *Clinical Rehabilitation* 1999;13(5):373-83.

Pandyan AD, Price CIM, Rodgers H, Barnes MP, Johnson GR. Biomechanical examination of a commonly used measure of spasticity. *Clinical Biomechanics* 2001;16(10):859-865.

Pandyan AD, Price CIM, Barnes MP, Johnson GR. A biomechanical investigation into the validity of the Modified Ashworth Scale as a measure of elbow spasticity. *Clinical Rehabilitation* 2003;17:290-294.

Patrick E, Ada L. The Tardieu Scale differentiates contracture from spasticity whereas the Ashworth Scale is confounded by it. *Clinical Rehabilitation* 2006;20(2):173-82.

Perell K, Scremin A, Scremin O, Kunkel C. Quantifying muscle tone in spinal cord injury patients using isokinetic dynamometric techniques. *Paraplegia* 1996;34(1):46-53.

Pisano F, Miscio G, Del Conte C, Pianca D, Candeloro E, Colombo R. Quantitative measures of spasticity in post-stroke patients. *Clinical Neurophysiology* 2000;111(6):1015-22.

Pizzi A, Carlucci G, Falsini C, Verdesca S, Grippo A. Evaluation of upper-limb spasticity after stroke: a clinical and neurophysiologic study. *Archives of Physical Medicine and Rehabilitation* 2005;86:410-415.

Platz T, Eickhof C, Nuyens G, Vuadens P. Clinical scales for the assessment of spasticity, associated phenomena, and function: a systematic review of the literature. *Disability and Rehabilitation* 2005Jan7-21;27(1/2):7-18.

Nuyens G, De Weerdt W, Ketelaer P, Feys H, De Wolf L, Hantson L, Nieuwboer A, Spaepen A, Carton H. Inter-rater reliability of the Ashworth scale in multiple sclerosis. *Clinical Rehabilitation* 1994;8:286-292.

Skold C. Spasticity in spinal cord injury: self- and clinically rated intrinsic fluctuations and intervention-induced changes. *Archives of Physical Medicine and Rehabilitation* 2000;81:144-149.

Skold C, Harms-Ringdahl K, Hultling C, Levi R, Seiger A. Simultaneous Ashworth measurements and electromyographic recordings in tetraplegic patients. *Archives of Physical Medicine and Rehabilitation* 1998;79:959-965.

Sloan RL, Sinclair E, Thompson J, Taylor S, Pentland B. Inter-rater reliability of the Modified Ashworth Scale for spasticity in hemiplegic patients. *International Journal of Rehabilitation Research* 1992;15:158-61.

Tederko P, Krasuski M, Czech J, Dargiel A, Garwacka-Jodzis I, Wojciechowska A. Reliability of clinical spasticity measurements in patients with cervical spinal cord injury. *Ortopedia Traumatologia Rehabilitacja* 2007;9(5):467-83.

Voerman GE, Gregoric M, Hermens HJ. Neurophysiological methods for the assessment of spasticity: the Hoffmann reflex, the tendon reflex, and the stretch reflex. *Disability and Rehabilitation* 2005;27(1/2):33-68.

Waninge A, Rook RA, Dijkhuizen A, Gielen E, van der Schans CP. Feasibility, test-retest reliability, and interrater reliability of the Modified Ashworth Scale and Modified Tardieu Scale in persons with profound intellectual and multiple disabilities. *Research in Developmental Disabilities* 2011;32(2):613-20.

Yam WKL, Leung MSM. Interrater reliability of Modified Ashworth Scale and Modified Tardieu Scale in children with spastic cerebral palsy. *Journal of Child Neurology* 2006;21(12):1031-5.