

## **Tardieu References:**

Abolhasani H, Ansari NN, Naghdi S, Mansouri K, Ghotbi N, Hasson S. Comparing the validity of the Modified Modified Ashworth Scale (MMAS) and the Modified Tardieu Scale (MTS) in the assessment of wrist flexor spasticity in patients with stroke: protocol for a neurophysiological study. *BMJ Open* 2012;Nov19:2(6):e001394.

Akpınar P, Atıcı A, Özkan FU, Aktas I, Kulcu DG, Sari A, Durmus B. Reliability of the Modified Ashworth Scale and Modified Tardieu Scale in patients with spinal cord injuries. *Spinal Cord* 2017;55(10):944-949.

Alhusaini AA, Dean CM, Crosbie J, Shepherd RB, Lewis J. Evaluation of spasticity in children with cerebral palsy using Ashworth and Tardieu Scales compared with laboratory measures. *Journal of Child Neurology* 2010;25(10):1242-7.

Ansari NN, Naghdi S, Hasson S, Azarsa MH, Azarnia S. The Modified Tardieu Scale for the measurement of elbow flexor spasticity in adult patients with hemiplegia. *Brain Injury* 2008;22(13-14):1007-1012.

Ansari NN, Naghdi S, Hasson S, Rastgoo M, Amini M, Forogh B. Clinical assessment of ankle plantarflexor spasticity in adult patients after stroke: inter- and intra-rater reliability of the Modified Tardieu Scale. *Brain Injury* 2013;27(5):605-612.

Bar-On L, Van Campenhout A, Desloovere K, Aertbeliën E, Huenaerts C, Vandendooren B, Molenaers G. Is an instrumented spasticity assessment an improvement over clinical spasticity scales in assessing and predicting the response to integrated botulinum toxin type a treatment in children with cerebral palsy? *Archives of Physical Medicine and Rehabilitation* 2014;95(3), 515-523.

Boyd RN, Barwood SA, Ballieu C, Graham HK. Validity of a clinical measure of spasticity in children with cerebral palsy in a double-blinded randomised controlled clinical trial. *Developmental Medicine and Child Neurology* 1998;40(suppl 78):7.

Boyd RN, Graham HK. Objective measurement of clinical findings in the use of botulinum toxin type A for the management of children with cerebral palsy. *European Journal of Neurology* 1999;6(suppl 4):S23-S35.

Fosang AL, Galea MP, McCoy AT, Reddihough DS. Measures of muscle and joint performance in the lower limb of children with cerebral palsy. *Developmental Medicine and Child Neurology* 2003;45(10):664-670.

Gholami S, Ansari NN, Naghdi S, Tabatabaei A, Jannat D, Senobari M, Dadgoo M. Biomechanical investigation of the modified Tardieu Scale in assessing knee extensor spasticity poststroke. *Physiotherapy Research International* 2018;23(1):e1698.

Gracies JM, Burke K, Clegg N, Browne R, Rushing C, Fehlings D, Matthews D, Tilton A, Delgado MR. Reliability of the Tardieu Scale for assessing spasticity in children with cerebral palsy. *Archives of Physical Medicine and Rehabilitation* 2010;91(3):421-428.

Gracies J-M, Marosszeky, J E, Renton R, Sandanam J, Gandevia SC, Burke D. Short-term effects of dynamic lycra splints on upper limb in hemiplegic patients. *Archives of Physical Medicine and Rehabilitation* 2000;81:1547-55.

Haugh AB, Pandyan AD, Johnson GR. A systematic review of the Tardieu Scale for the measurement of spasticity. *Disability and Rehabilitation* 2006;28(15):899-907.

Held JP, Peierrot-Deseilligny E. *Reeducation motrice des affections neurologiques*. Paris: JB Bailliere et fils, 1969:31-42.

Li F, Wu Y, Li X. Test-retest reliability and inter-rater reliability of the Modified Tardieu Scale and the Modified Ashworth Scale in hemiplegic patients with stroke. *European Journal of Physical and Rehabilitation Medicine* 2014;50(1):9-15.

Mackey AH, Walt SE, Lobb G, Stott NS. Intraobserver reliability of the modified Tardieu scale in the upper limb of children with hemiplegia. *Developmental Medicine and Child Neurology* 2004;46(4):267-72.

Mehrholtz J, Wagner K, Meißner D, Grundmann K, Zange C, Koch R, Pohl M. Reliability of the Modified Tardieu Scale and the Modified Ashworth Scale in adult patients with severe brain injury: a comparison study. *Clinical Rehabilitation* 2005;19(7):751-9.

Morris SL, Williams G. A historical review of the evolution of the Tardieu Scale. *Brain Injury* 2018;32:5:665-669.

Naghdi S, Ansari NN, Ghorbani-rad S, Senobari M, Sahraian MA. Intra-rater reliability of the Modified Tardieu Scale in patients with multiple sclerosis. *Neurological Sciences* 2017 Jan;38(1):93-99.

Numanoğlu A, Günel MK. Intraobserver reliability of Modified Ashworth Scale and Modified Tardieu Scale in the assessment of spasticity in children with cerebral palsy. *Acta Orthopaedica et Traumatologica Turc* 2012;46(3):196-200.

Patrick E, Ada L. The Tardieu Scale differentiates contracture from spasticity whereas the Ashworth Scale is confounded by it. *Clinical Rehabilitation* 2006;20(2):173-82.

Paulis WD, Horemans HL, Brouwer BS, Stam HJ. Excellent test-retest and inter-rater reliability for Tardieu Scale measurements with inertial sensors in elbow flexors of stroke patients. *Gait and Posture* 2011;33(2): 185-189.

Scholtes VAB, Becher JG, Beelen A, Lankhorst GJ. Clinical assessment of spasticity in children with cerebral palsy: a critical review of available instruments. *Developmental Medicine and Child Neurology* 2006;48(1), 64-73.

Singh P, Joshua AM, Ganesham S, Suresh S. Intra-rater reliability of the modified Tardieu scale to quantify spasticity in elbow flexors and ankle plantar flexors in adult stroke subjects. *Annals of Indian Academy of Neurology* 2011;14(1): 23-26.

Tardieu G, Shentoub S, Delarue R. A la recherche d'une technique de mesure de la spasticite. *Rev Neurol (Paris)* 1954;91:143-144.

Waninge A, Rook RA, Dijkhuizen A, Gielen E, van der Schans CP. Feasibility, test-retest reliability, and interrater reliability of the Modified Ashworth Scale and Modified Tardieu Scale in persons with profound intellectual and multiple disabilities. *Research in Developmental Disabilities* 2011;32(2):613-20.

Yam WKL, Leung MSM. Interrater reliability of Modified Ashworth Scale and Modified Tardieu Scale in children with spastic cerebral palsy. *Journal of Child Neurology* 2006;21(12):1031-5.